

Atty. Dkt. No. 025098-0701

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

1. (Previously presented) A method for designing a specific polyamide



wherein

X_1 , X_2 , X_m , $X_{(m+1)}$, $X_{(2m-1)}$, and X_{2m} are carboxamide residues forming carboxamide binding pairs X_1/X_{2m} , $X_2/X_{(2m-1)}$, $X_m/X_{(m+1)}$;

γ is γ -aminobutyric acid or 2,4 diaminobutyric acid, and

R_1 is $-\text{NH}(\text{CH}_2)_{0-100}\text{NR}_2\text{R}_3$, $-\text{NH}(\text{CH}_2)_{0-12}\text{CONH}(\text{CH}_2)_{0-100}\text{NR}_2\text{R}_3$, or $-\text{NHR}_2$, where R_2 and R_3 are independently selected from the group consisting of H, Cl, NO, N-acetyl, benzyl, C_{1-100} alkyl, C_{1-100} alkylamine, C_{1-100} alkyldiamine, C_{1-100} alkylcarboxylate, C_{1-100} alkenyl, a C_{1-100} alkynyl, and C_{1-100} alkyl-L, where L is selected from the group consisting of arylboronic acids, biotins, polyhistidines comprised from about 2 to 8 amino acids, haptens, solid phase supports, oligodeoxynucleotides, N-ethylnitrosourea, fluorescein, bromoacetamide, iodoacetamide, DL- α -lipoic acid, acridine, captothesin, pyrene, mitomycin, texas red, anthracene, anthrinilic acid, avidin, DAPI, and oligodeoxynucleotide, isosulfan blue, malachite green, psoralen, ethyl red, 4-(psoraen-8-yloxy)-butyrate, tartaric acid, and (+)- α -tocopheral, suitable for use as a DNA-binding ligand that is selective for identified target DNA-sequences $5'-\text{WN}_1\text{N}_2 \dots \text{N}_m\text{W}-3'$ where m is an integer having a value from 3 to 6, the method comprising:

Atty. Dkt. No. 025098-0701

(a) identifying a target sequence of double stranded DNA having the form 5'-WN₁N₂ ... N_mW-3', N₁N₂ ... N_m being the sequence to be bound by carboxamide residues, wherein each N is independently chosen from the group A, G, C, and T, each W is independently chosen from the group A and T, and m is an integer having a value from 3 to 6;

(b) representing the identified sequence as 5'-Wab ... xW-3', wherein a is a first nucleotide to be bound by the X₁ carboxamide residue, b is a second nucleotide to be bound by the X₂ carboxamide residue, and x is the corresponding nucleotide to be bound by the X_m carboxamide residue;

(c) defining a as A, G, C, or T to correspond to the first nucleotide to be bound by a carboxamide residue in the identified sequence;

(d) selecting Im as the X₁ carboxamide residue and Py as the X_{2m} carboxamide residue if a = G;

(e) selecting Py as the X₁ carboxamide residue and Im as the X_{2m} carboxamide residue if a = C;

(f) selecting Hp as the X₁ carboxamide residue and Py as the X_{2m} carboxamide residue if a = T;

(g) selecting Py as the X₁ carboxamide residue and Hp as the X_{2m} carboxamide residue if a = A; and

(h) repeating steps c - g for b through x until all carboxamide residues are selected;

Atty. Dkt. No. 025098-0701

wherein Im is N-methylimidazole, Hp is 3-hydroxy-N-methylpyrrole, Py is N-methylpyrrole, A is adenine, G is guanine, C is cytosine, and T is thymine; and synthesizing the polyamide.

2. (Cancelled)

3. (Previously presented) The method of claim 1 further comprising the step of determining if the binding affinity of the polyamide to the identified target sequence is subnanomolar.

4. (Previously presented) The method of claim 1 further comprising the step of determining if the polyamide exhibits a binding affinity that is at least ten-fold higher for said identified target sequence compared to a non-target DNA sequence.

5. (Previously presented) The method of claim 1 further comprising the step of replacing at least one pyrrole residue with a β -alanine residue.

6-41. (Cancelled)

42. (Previously presented) The method of claim 1 wherein the identified target DNA sequence is a regulatory sequence.

43. (Previously presented) The method of claim 1 wherein the identified target DNA sequence is a promoter sequence.

44. (Previously presented) The method of claim 1 wherein the identified target DNA sequence is a coding sequence.

Atty. Dkt. No. 025098-0701


45. (Previously presented) The method of claim 1 wherein the identified target DNA sequence is a non-coding sequence.

46. (Previously presented) A polyamide composition produced by the method of claim 1 wherein the binding of the carboxamide binding pairs to the identified target DNA sequence modulates the expression of a gene.

47. (Previously presented) A composition comprising an effective amount of a polyamide produced by the method of claim 1 and a pharmacologically suitable excipient.

48. (Previously presented) A diagnostic kit comprising a polyamide produced by the method of claim 1.

49. (Previously presented) A polyamide designed by the method of claim 1, having the structure:



wherein

R₄ is selected from the group consisting of H, NH₂, SH, Cl, Br, F, N-acetyl, and N-formyl;

R₅ is H or NH₂;

Atty. Dkt. No. 025098-0701

R_1 is $-NH(CH_2)_{0-100}NR_2R_3$, $-NH(CH_2)_{0-12}CONH(CH_2)_{0-100}NR_2R_3$, or $-NHR_2$, where R_2 and R_3 are independently selected from the group consisting of H, Cl, NO, N-acetyl, benzyl, C_{1-100} alkyl, C_{1-100} alkylamine, C_{1-100} alkyldiamine, C_{1-100} alkylcarboxylate, C_{1-100} alkenyl, a C_{1-100} alkynyl, and C_{1-100} alkyl-L, where L is selected from the group consisting of arylboronic acids, biotins, polyhistidines comprised from about 2 to 8 amino acids, haptens, solid phase supports, oligodeoxynucleotides, N-ethylnitrosourea, fluorescein, bromoacetamide, iodoacetamide, DL- α -lipoic acid, acridine, captothesin, pyrene, mitomycin, texas red, anthracene, anthranilic acid, avidin, DAPI, and oligodeoxynucleotide, isosulfan blue, malachite green, psoralen, ethyl red, 4-(psoralen-8-yloxy)-butyrate, tartaric acid, and (+)- α -tocopheral;

each X is independently selected from the group consisting of N, CH, and COH;

each a is an integer from 2 to 5; and

each b is an integer from 3 to 6.

50. (New) A method for designing a specific polyamide



wherein

X_1 , X_2 , X_m , $X_{(m+1)}$, $X_{(2m-1)}$, and X_{2m} are carboxamide residues forming carboxamide binding pairs X_1/X_{2m} , $X_2/X_{(2m-1)}$, $X_m/X_{(m+1)}$, and wherein one carboxamide binding pair is substituted with a β/β , wherein β is β -alanine;

γ is γ -aminobutyric acid or 2,4 diaminobutyric acid, and

Atty. Dkt. No. 025098-0701

R_1 is $-\text{NH}(\text{CH}_2)_{0-100}\text{NR}_2\text{R}_3$, $-\text{NH}(\text{CH}_2)_{0-12}\text{CONH}(\text{CH}_2)_{0-100}\text{NR}_2\text{R}_3$, or $-\text{NHR}_2$, where R_2 and R_3 are independently selected from the group consisting of H, Cl, NO, N-acetyl, benzyl, C_{1-100} alkyl, C_{1-100} alkylamine, C_{1-100} alkyldiamine, C_{1-100} alkylcarboxylate, C_{1-100} alkenyl, a C_{1-100} alkynyl, and C_{1-100} alkyl-L, where L is selected from the group consisting of arylboronic acids, biotins, polyhistidines comprised from about 2 to 8 amino acids, haptens, solid phase supports, oligodeoxynucleotides, N-ethylnitrosourea, fluorescein, bromoacetamide, iodoacetamide, DL- α -lipoic acid, acridine, captothesin, pyrene, mitomycin, texas red, anthracene, anthrinilic acid, avidin, DAPI, and oligodeoxynucleotide, isosulfan blue, malachite green, psoralen, ethyl red, 4-(psoraen-8-yloxy)-butyrate, tartaric acid, and (+)- α -tocopheral, suitable for use as a DNA-binding ligand that is selective for identified target DNA-sequences $5' \text{--} \text{WN}_1\text{N}_2 \dots \text{N}_m\text{W} \text{--} 3'$ where m is an integer having a value from 3 to 6, the method comprising:

(a) identifying a target sequence of double stranded DNA having the form $5' \text{--} \text{WN}_1\text{N}_2 \dots \text{N}_m\text{W} \text{--} 3'$, $\text{N}_1\text{N}_2 \dots \text{N}_m$ being the sequence to be bound by carboxamide residues, wherein each N is independently chosen from the group A, G, C, and T, each W is independently chosen from the group A and T, and m is an integer having a value from 3 to 6;

(b) representing the identified sequence as $5' \text{--} \text{Wab} \dots \text{xW} \text{--} 3'$, wherein a is a first nucleotide to be bound by the X_1 carboxamide residue, b is a second nucleotide to be bound by the X_2 carboxamide residue, and x is the corresponding nucleotide to be bound by the X_m carboxamide residue;

(c) defining a as A, G, C, or T to correspond to the first nucleotide to be bound by a carboxamide residue in the identified sequence;

Atty. Dkt. No. 025098-0701

(d) selecting Im as the X_1 carboxamide residue and Py as the X_{2m} carboxamide residue if $a = G$;

(e) selecting Py as the X_1 carboxamide residue and Im as the X_{2m} carboxamide residue if $a = C$;

(f) selecting Hp as the X_1 carboxamide residue and Py as the X_{2m} carboxamide residue if $a = T$;

(g) selecting Py as the X_1 carboxamide residue and Hp as the X_{2m} carboxamide residue if $a = A$; and

(h) repeating steps c – g for b through x until all carboxamide residues are selected;

wherein Im is N-methylimidazole, Hp is 3-hydroxy-N-methylpyrrole, Py is N-methylpyrrole, A is adenine, G is guanine, C is cytosine, and T is thymine; and

synthesizing the polyamide.